Tendon Disorders



What are tendons?

Tendons are rope-like bundles or bands of strong, smooth, shiny fibres that attach muscles to bones. Tendons transfer force from the muscle to the bone to produce the movement of joints. There are two types of tendons: tendons without sheaths (Figure 1), and tendons with sheaths (Figure 2).

Figure 1

Figure 2

What are tendon disorders?

Tendon disorders, or tendinopathies, are medical conditions that result in the tendons not functioning normally. Tendinitis is a disorder of tendons without sheaths and tenosynovitis is a disorder of tendons with sheaths.

What is tendinitis?

Traditionally, tendinitis (also called tendonitis) is the term used to describe an inflammation or irritation of a tendon. With repetitive or prolonged activities,

forceful exertion, awkward and static postures, vibration, and localized mechanical stress, the tendons fibers can tear apart in much the same way a rope becomes frayed.

These tendon changes trigger an inflammatory response. Inflammation is a localized response of tissue to injury. Over time, inflamed tendons become thickened, bumpy, and irregular. Without rest and time for the tissue to heal, tendons can become permanently weakened.

Because the term "tendonitis" implies inflammation is present, this term is rarely used. "Tendinopathy" or "tendinosis" is the preferred term to describe chronic pain associated with a symptomatic tendon.

What is tenosynovitis?

Tenosynovitis is an inflammation of the tendon sheath. The inner walls of the sheath produce a slippery fluid, called synovial fluid, which lubricates the tendon. With repetitive or prolonged activities, forceful exertion, awkward and static posture, vibration, and localized mechanical stress, the lubrication system may malfunction. It may not produce enough fluid, or it may produce a fluid with poor lubricating qualities. Friction between the tendon and its sheath results when the lubricating system fails. This leads to inflammation and swelling of the tendon sheath. Repeated episodes of inflammation cause fibrous tissue to form that thickens the tendon sheath and hinders tendon movement.

Where do tendon disorders occur?

The most common sites for occupational tendon disorders are (Figure 3):

Figure 3

Shoulder

- Bicipital tendinopathy disorder of the tendon to the biceps.
- Rotator cuff tendinopathy disorder of the tendons that rotate the humerus [upper arm bone] and help raise the arm.

Elbow

Lateral epicondylitis or tennis elbow — inflammation of the tendon that attaches some

forearm muscles to a "bump" or projection [lateral epicondyle] on the side of the humerus, just above the elbow.

Hand and Wrist

- Flexor tendinopathy disorder of the tendons in the palm side of the wrist and hand
- Extensor tendinopathy disorder of the tendons in the back of the hand and wrist.
- Flexor tenosynovitis inflammation of the tendon sheaths in the palm side of the wrist and hand.
- Extensor tenosynovitis inflammation of the tendons in the back of the hand and wrist.
- De Quervain's disease inflammation of the tendon sheaths at the base of the thumb.
- Dupuytren's contracture in this condition, the tissues (fascia) under the skin on the palm of the hand thicken and shorten enough that the tendons connected to the fingers cannot move freely; this leads to a non-inflammatory degeneration of the tendons of the palmaris muscle in the hand.

Shoulder tendinopathy

The most common tendon disorders of the shoulder are bicipital tendinopathy and rotator cuff tendinopathy. Bicipital tendinopathy may be caused by inflammation of the tendon around the biceps muscle, or by degeneration of the tendon from repetitive overhead motion or from the normal aging process. Rotator cuff tendinopathy is also known as supraspinatus tendinitis, subdeltoid bursitis, subacromial bursitis or partial tear of the rotator cuff.

Tendon disorders of the shoulder are seen among workers exposed to high repetitive movements requiring a significant use of force. Workers doing jobs requiring awkward postures such as overhead work, arm elevation, and specific postures relative to the degree of arm bending and arm elevation.

Elbow tendinopathy

Tendinopathy of the elbow also known as tennis elbow or lateral epicondylitis is associated with jobs that require repeated or forceful movements of the fingers, wrist, and forearm. Specific movements associated with the development of tendinopathy of the elbow include simultaneous rotation of the forearm and bending of the wrist; stressful gripping of objects with inward or outward movement of the forearm; jerky, throwing motions.

Hand and wrist tendinopathy

Tendinopathy of the hand and wrist include a variety of diagnoses such as tenosynovitis, tendinitis, De Quervain's disease, and Dupuytren's contracture.

High risk occupation and activities associated with hand and wrist tendinitis include assembly line work, meat processing, manufacturing, knitting, typing, and piano playing.

There is some evidence that workers exposed to risk factors such as high force and high repetition are at increased risk for hand and wrist tendinopathy. Excessive texting on cellular phones has been found to be a potential risk factor for De Ouervain's disease from overuse of the thumbs.

Patellar (knee) tendinitis

Patellar tendinitis is an injury to the tendon connecting your kneecap (patella) to your shinbone. This tendon helps to extend your knee so you can kick, run and jump. Running and jumping are the most common causes, but sudden increases in how hard or

how often you run or jump can add stress. While common in athletes, people in occupations with similar movements may be at risk.

What are the symptoms?

The clinical presentation of tendon disorders is characterized by the presence of pain on the site of injury. Often, there is a decrease in strength or functional ability as well. Specific physical findings on examination include tenderness when the area over the affected tendon is touched and may be associated with swelling, redness, and restriction of movement.

What causes tendon disorders?

Often it is a combination of factors that leads to a tendon disorder, and every situation is unique. Causes can include overuse as well as age, injury, or disease related changes in the tendon. Risk factors for tendon disorders can include excessive force, repetitive movements, frequent overhead reaching, vibration, and awkward postures.

What is the treatment?

Treatment of tendon disorders is of a conservative nature and rarely requires surgery.

Avoiding those activities that caused or aggravated the disorder will prevent the condition from persisting or recurring. Pain relievers, corticosteroids or plateletrich plasma may be options to discuss with your medical professional. Physical therapy can help stretch and strengthen the affected muscles and tendons.

How can we prevent tendon disorders?

Tendon disorders have been associated with repetitive or prolonged activities, forceful exertion, awkward and static posture, vibration, and localized mechanical stress. The prevention of tendon disorders should include identifying and remedying these risk factors

Good design of work practices and equipment should aim at reducing repetitive movements, awkward postures and static posture (periods spent in one position). Job design should also aim at minimizing the need to use forceful exertion and making sure that rest and work breaks are properly used.

The prevention of tendon disorders should also include training and education. To be successful a training and education program must be organized, consistent and ongoing. Everyone at work, including workers, managers, health and safety representatives, and so on, must get actively involved. Education and training can include how to use the right technique during activity or work to lessen the possible issues (e.g., stretch, strengthen muscles, appropriate lifting techniques, good ergonomic set-up, etc.)

Source: © Copyright 1997-2021 CCOHS